

University of Wollongong

Research Online

Faculty of Health and Behavioural Sciences -
Papers (Archive)

Faculty of Science, Medicine and Health

2007

The hollow face illusion in infancy

E Nakato

Chuo University

Harold C. Hill

University of Wollongong, harry@uow.edu.au

Y Otsuka

Tokyo Women's Medial College

S Kanazawa

Shukutoku University

M Yamaguchi

Chuo University

Follow this and additional works at: <https://ro.uow.edu.au/hbspapers>



Part of the [Arts and Humanities Commons](#), [Life Sciences Commons](#), [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Nakato, E; Hill, Harold C.; Otsuka, Y; Kanazawa, S; and Yamaguchi, M: The hollow face illusion in infancy 2007, 150-150.
<https://ro.uow.edu.au/hbspapers/982>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

The hollow face illusion in infancy

Abstract

Abstract presented at The 30th European Conference on Visual Perception, 27-31 August 2007, Arezzo, Italy

Keywords

infancy, illusion, face, hollow

Disciplines

Arts and Humanities | Life Sciences | Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Nakato, E., Hill, H. C., Otsuka, Y., Kanazawa, S. & Yamaguchi, M. (2007). The hollow face illusion in infancy. *Perception*, 36 (Supplement), 150.

Thursday

boundary or the ability to discriminate within-category and between-category pairs of faces. In contrast, for the sadness–fear continuum, older adults differed from younger adults in the locus of the category boundary and were less accurate in discriminating between-category but not within-category pairs of faces. This suggests that difficulties in processing emotional expressions of fear and sadness may be caused by age-related changes in sensitivity to variation between perceptual categories.

[Supported by a University of Otago Research Grant.]

◆ **Does visual categorization really shape feature selectivity in the primate temporal cortex?**

YNajian, Sh Talehy Moineddin¶, F Sodagari¶ (Department of Basic Sciences, Nikan Health Research Institute, Tehran, Iran; ¶ Iran University of Medical Sciences, Tehran, Iran; e-mail: yousef.najian@gmail.com)

Activity in the human temporal cortex is thought to be sensitive to the categorization level of the stimuli and to depend on the expertise of the observer. Sigala and Logothetis (2002 *Nature* **415** 318–320) published an article with the aim of testing whether inferior temporal cortex neurons respond selectively to object features that constitute the relevant dimensions for visual object categorization. They defined parameterized line drawings of faces each consisting of an outline and four varying features as stimuli and trained two monkeys in a categorization task. They claimed that the categories were separable along two of the four dimensions of the stimuli. On examining their stimuli again, we found that the face categories were separable along the combination of one feature with each of the remaining features. Therefore, there was only one diagnostic dimension and the monkey could optionally select the second one for categorization. Assuming that the selectivity index for the real unitary diagnostic feature was much larger than for the other three features, we find the interpretation to be different.

◆ **The hollow face illusion in infancy**

E Nakato, H Hill¶, Y Otsuka§, S Kanazawa#, M Yamaguchi (Department of Psychology, Chuo University, Hachioji, Japan; ¶ University of Wollongong, Wollongong, Australia; § Tokyo Women's Medical University, Tokyo, Japan; # Shukutoku University, Chiba, Japan; e-mail: enakato@komazawa-u.ac.jp)

The hollow-face illusion is perceived on the basis of two assumptions: (i) top–down knowledge for faces, (ii) convexity preference of patterns of shading (Hill and Johnson, 2007 *Perception* 199–223). We examined the effect of the hollow-face illusion in infants aged 6–8 months. Infants were first familiarized for 80 s with a male face either in the convex or the concave condition. In both conditions each face was moved from side to side over 20°. After familiarization, infants were tested with a pair of novel convex and concave faces in two 10 s trials. If infants perceived a hollow mask as a convex face, they would show a novelty preference for the novel concave faces in test phase. The results indicated infants' preference for concave faces only in the convex condition. The results suggest that the hollow-face illusion did not appear even in 8-month-old infants.

[Supported by RISTEX and PRESTO (JST) and by a Grant-in-Aid for Scientific Research (18000090) from the JSPS.]

◆ **The effects of camera and light angles on the recognition of emotion from the face**

M Oda (Department of Psychology, Ritsumeikan University, Kyoto, Japan; e-mail: oda@lt.ritsumei.ac.jp)

The effects of camera and light angles on the recognition of facial expressions were investigated. Facial expressions and neutral faces were taken by five cameras simultaneously under four light direction conditions. The participants rated emotional intensity and reality of the stimuli. Angry faces were rated stronger in the front camera and in the front light, and weaker in the side camera and in the side light. Smiling faces were regarded as weaker in the upper camera and in the upper/side light. On the other hand, neutral faces were regarded as angry faces in the diagonal/side camera condition and upper light condition, and as smiling faces under the low light condition. These results indicate that the camera and light angles influence the facial impression, and intensity of emotion changes by the combination of emotional types, camera angles, and light angles. Moreover, the effects of camera and light angles on the neutral face and emotional expressions face could be caused by different mechanisms respectively.

◆ **The effects of adaptation duration on contrast thresholds in face identification**

I Oruc, J J Barton (Department of Ophthalmology and Visual Sciences, University of British Columbia, Vancouver, Canada; e-mail: ipek@psych.ubc.ca)

We report a novel face-adaptation paradigm in which we probe the effect of prior exposure on contrast thresholds for face identification. Each trial consisted of an adaptation period (showing a